

2011

Models for Discrete Quantum Gravity

S. Gudder

Follow this and additional works at: https://digitalcommons.du.edu/math_preprints



Part of the [Mathematics Commons](#)

Recommended Citation

Gudder, S., "Models for Discrete Quantum Gravity" (2011). *Mathematics Preprint Series*. 35.
https://digitalcommons.du.edu/math_preprints/35

This Article is brought to you for free and open access by the Department of Mathematics at Digital Commons @ DU. It has been accepted for inclusion in Mathematics Preprint Series by an authorized administrator of Digital Commons @ DU. For more information, please contact jennifer.cox@du.edu.

MODELS FOR DISCRETE QUANTUM GRAVITY

S. Gudder

Department of Mathematics

University of Denver

Denver, Colorado 80208, U.S.A.

sgudder@du.edu

Abstract

We first discuss a framework for discrete quantum processes (DQP). It is shown that the set of q -probability operators is convex and its set of extreme elements is found. The property of consistency for a DQP is studied and the quadratic algebra of suitable sets is introduced. A classical sequential growth process is “quantized” to obtain a model for discrete quantum gravity called a quantum sequential growth process (QSGP). Two methods for constructing concrete examples of QSGP are provided.

1 Introduction

In a previous article, the author introduced a general framework for a discrete quantum gravity [3]. However, we did not include any concrete examples or models for this framework. In particular, we did not consider the problem of whether nontrivial models for a discrete quantum gravity actually exist. In this paper we provide a method for constructing an infinite number of such models. We first make a slight modification of our definition of a discrete quantum process (DQP) ρ_n , $n = 1, 2, \dots$. Instead of requiring that ρ_n be a state on a Hilbert space H_n , we require that ρ_n be a q -probability operator on H_n . This latter condition seems more appropriate from a probabilistic viewpoint and instead of requiring $\text{tr}(\rho_n) = 1$, this condition normalizes the

corresponding quantum measure. By superimposing a concrete DQP on a classical sequential growth process we obtain a model for discrete quantum gravity that we call a quantum sequential growth process.

Section 2 considers the DQP formalism. We show that the set of q -probability operators is a convex set and find its set of extreme elements. We discuss the property of consistency for a DQP and introduce the so-called quadratic algebra of suitable sets. The suitable sets are those on which well-defined quantum measures (or quantum probabilities) exist.

Section 3 reviews the concept of a classical sequential growth process (CSGP) [1, 4, 5, 6, 8, 9]. The important notions of paths and cylinder sets are discussed. In Section 4 we show how to “quantize” a CSGP to obtain a quantum sequential growth process (QSGP). Some results concerning the consistency of a DQP are given. Finally, Section 5 provides two methods for constructing examples of QSGP.

2 Discrete Quantum Processes

Let $(\Omega, \mathcal{A}, \nu)$ be a probability space and let

$$H = L_2(\Omega, \mathcal{A}, \nu) = \left\{ f: \Omega \rightarrow \mathbb{C}, \int |f|^2 d\nu < \infty \right\}$$

be the corresponding Hilbert space. Let $\mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots \subseteq \mathcal{A}$ be an increasing sequence of sub σ -algebras of \mathcal{A} that generate \mathcal{A} and let $\nu_n = \nu | \mathcal{A}_n$ be the restriction of ν to \mathcal{A}_n , $n = 1, 2, \dots$. Then $H_n = L_2(\Omega, \mathcal{A}_n, \nu_n)$ forms an increasing sequence of closed subspaces of H called a *filtration* of H . A bounded operator T on H_n will also be considered as a bounded operator on H by defining $Tf = 0$ for all $f \in H_n^\perp$. We denote the characteristic function χ_Ω of Ω by 1. Of course, $\|1\| = 1$ and $\langle 1, f \rangle = \int f d\nu$ for every $f \in H$. A *q-probability operator* is a bounded positive operator ρ on H that satisfies $\langle \rho 1, 1 \rangle = 1$. Denote the set of q -probability operators on H and H_n by $\mathcal{Q}(H)$ and $\mathcal{Q}(H_n)$, respectively. Since $1 \in H_n$, if $\rho \in \mathcal{Q}(H_n)$ by our previous convention, $\rho \in \mathcal{Q}(H)$. Notice that a positive operator $\rho \in \mathcal{Q}(H)$ if and only if $\|\rho^{1/2} 1\| = 1$ where $\rho^{1/2}$ is the unique positive square root of ρ .

A rank 1 element of $\mathcal{Q}(H)$ is called a *pure q-probability operator*. Thus $\rho \in \mathcal{Q}(H)$ is pure if and only if ρ has the form $\rho = |\psi\rangle\langle\psi|$ for some $\psi \in H$ satisfying

$$|\langle 1, \psi \rangle| = \left| \int \psi d\nu \right| = 1$$

We then call ψ a q -probability vector and we denote the set of q -probability vectors by $\mathcal{V}(H)$ and the set of pure q -probability operators by $\mathcal{Q}_p(H)$. Notice that if $\psi \in \mathcal{V}(H)$, then $\|\psi\| \geq 1$ and $\|\psi\| = 1$ if and only if $\psi = \alpha 1$ for some $\alpha \in \mathbb{C}$ with $|\alpha| = 1$. Two operators $\rho_1, \rho_2 \in \mathcal{Q}(H)$ are *orthogonal* if $\rho_1 \rho_2 = 0$.

Theorem 2.1. (i) $\mathcal{Q}(A)$ is a convex set and $\mathcal{Q}_p(H)$ is its set of extreme elements. (ii) $\rho \in \mathcal{Q}(H)$ is of trace class if and only if there exists a sequence of mutually orthogonal $\rho_i \in \mathcal{Q}_p(H)$ and $\alpha_i > 0$ with $\sum \alpha_i = 1$ such that $\rho = \sum \alpha_i \rho_i$ in the strong operator topology. The ρ_i are unique if and only if the corresponding α_i are distinct.

Proof. (i) If $0 < \lambda < 1$ and $\rho_1, \rho_2 \in \mathcal{Q}(H)$, then $\rho = \lambda \rho_1 + (1 - \lambda) \rho_2$ is a positive operator and

$$\langle \rho 1, 1 \rangle = \langle (\lambda \rho_1 + (1 - \lambda) \rho_2) 1, 1 \rangle = \lambda \langle \rho_1 1, 1 \rangle + (1 - \lambda) \langle \rho_2 1, 1 \rangle = 1$$

Hence, $\rho \in \mathcal{Q}(H)$ so $\mathcal{Q}(H)$ is a convex set. Suppose $\rho \in \mathcal{Q}_p(H)$ and $\rho = \lambda \rho_1 + (1 - \lambda) \rho_2$ where $0 < \lambda < 1$ and $\rho_1, \rho_2 \in \mathcal{Q}(H)$. If $\rho_1 \neq \rho_2$, then $\text{rank}(\rho) \neq 1$ which is a contradiction. Hence, $\rho_1 = \rho_2$ so ρ is an extreme element of $\mathcal{Q}(H)$. Conversely, suppose $\rho \in \mathcal{Q}(H)$ is an extreme element. If the cardinality of the spectrum $|\sigma(\rho)| > 1$, then by the spectral theorem $\rho = \rho_1 + \rho_2$ where $\rho_1, \rho_2 \neq 0$ are positive and $\rho_1 \neq \alpha \rho_2$ for $\alpha \in \mathbb{C}$. If $\rho_1 1, \rho_2 1 \neq 0$, then $\langle \rho_1 1, 1 \rangle, \langle \rho_2 1, 1 \rangle \neq 0$ and we can write

$$\rho = \langle \rho_1 1, 1 \rangle \frac{\rho_1}{\langle \rho_1 1, 1 \rangle} + \langle \rho_2 1, 1 \rangle \frac{\rho_2}{\langle \rho_2 1, 1 \rangle}$$

Now $\langle \rho_1 1, 1 \rangle^{-1} \rho_1, \langle \rho_2 1, 1 \rangle^{-1} \rho_2 \in \mathcal{Q}(H)$ and

$$\langle \rho_1 1, 1 \rangle + \langle \rho_2 1, 1 \rangle = \langle \rho 1, 1 \rangle = 1$$

which is a contradiction. Hence, $\rho_1 1 = 0$ or $\rho_2 1 = 0$. Without loss of generality suppose that $\rho_2 1 = 0$. We can now write

$$\rho = \frac{1}{2} \rho_1 + \frac{1}{2} (\rho_1 + 2 \rho_2)$$

Now $\rho_1 1 \neq 0$, $(\rho_1 + 2 \rho_2) 1 \neq 0$ and as before we get a contradiction. We conclude that $|\sigma(\rho)| = 1$. Hence, $\rho = \alpha P$ where P is a projection and $\alpha > 0$. If $\text{rank}(P) > 1$, then $P = P_1 + P_2$ where P_1 and P_2 are orthogonal nonzero projections so $\rho = \alpha P_1 + \alpha P_2$. Proceeding as before we obtain a contradiction. Hence, $\text{rank}(P) = 1$ so $\rho = \alpha P$ is pure. (ii) This follows from the spectral theorem. \square

Let $\{H_n: n = 1, 2, \dots\}$ be a filtration of H and let $\rho_n \in \mathcal{Q}(H_n)$, $n = 1, 2, \dots$. The n -decoherence functional $D_n: \mathcal{A}_n \times \mathcal{A}_n \rightarrow \mathbb{C}$ defined by

$$D_n(A, B) = \langle \rho_n \chi_B, \chi_A \rangle$$

gives a measure of the interference between A and B when the system is described by ρ_n . It is clear that $D_n(\Omega_n, \Omega_n) = 1$, $D_n(A, B) = \overline{D_n(B, A)}$ and $A \mapsto D_n(A, B)$ is a complex measure for all $B \in \mathcal{A}_n$. It is also well-known that if $A_1, \dots, A_r \in \mathcal{A}_n$ then the matrix with entries $D_n(A_j, A_k)$ is positive semidefinite. We define the map $\mu_n: \mathcal{A}_n \rightarrow \mathbb{R}^+$ by

$$\mu_n(A) = D_n(A, A) = \langle \rho_n \chi_A, \chi_A \rangle$$

Notice that $\mu_n(\Omega_n) = 1$. Although μ_n is not additive, it does satisfy the *grade-2 additivity condition*: if $A, B, C \in \mathcal{A}_n$ are mutually disjoint, then

$$\begin{aligned} \mu_n(A \cup B \cup C) &= \mu_n(A \cup B) + \mu_n(A \cup C) + \mu_n(B \cup C) \\ &\quad - \mu_n(A) - \mu_n(B) - \mu_n(C) \end{aligned} \quad (2.1)$$

We say that ρ_{n+1} is *consistent* with ρ_n if $D_{n+1}(A, B) = D_n(A, B)$ for all $A, B \in \mathcal{A}_n$. We call the sequence ρ_n , $n = 1, 2, \dots$, *consistent* if ρ_{n+1} is consistent with ρ_n for $n = 1, 2, \dots$. Of course, if the sequence ρ_n , $n = 1, 2, \dots$, is consistent, then $\mu_{n+1}(A) = \mu_n(A) \forall A \in \mathcal{A}_n$, $n = 1, 2, \dots$. A *discrete quantum process* (DQP) is a consistent sequence $\rho_n \in \mathcal{Q}(H_n)$ for a filtration H_n , $n = 1, 2, \dots$. A DQP ρ_n is *pure* if $\rho_n \in \mathcal{Q}_p(H_n)$, $n = 1, 2, \dots$.

If ρ_n is a DQP, then the corresponding maps $\mu_n: \mathcal{A}_n \rightarrow \mathbb{R}^+$ have the form

$$\mu_n(A) = \langle \rho_n \chi_A, \chi_A \rangle = \|\rho_n^{1/2} \chi_A\|^2$$

Now $A \mapsto \rho_n^{1/2} \chi_A$ is a vector-valued measure on \mathcal{A}_n . We conclude that μ_n is the squared norm of a vector-valued measure. In particular, if $\rho_n = |\psi_n\rangle\langle\psi_n|$ is a pure DQP, then $\mu_n(A) = |\langle\psi_n, \chi_A\rangle|^2$ so μ_n is the squared modulus of the complex-valued measure $A \mapsto \langle\psi_n, \chi_A\rangle$.

For a DQP $\rho_n \in \mathcal{Q}(H_n)$, we say that a set $A \in \mathcal{A}$ is *suitable* if $\lim \langle \rho_j \chi_A, \chi_A \rangle$ exists and is finite and in this case we define $\mu(A)$ to be the limit. We denote the set of suitable sets by $\mathcal{S}(\rho_n)$. If $A \in \mathcal{A}_n$ then

$$\lim \langle \rho_j \chi_A, \chi_A \rangle = \langle \rho_n \chi_A, \chi_A \rangle$$

so $A \in \mathcal{S}(\rho_n)$ and $\mu(A) = \mu_n(A)$. This shows that the algebra $\mathcal{A}_0 = \cup \mathcal{A}_n \subseteq \mathcal{S}(\rho_n)$. In particular, $\Omega \in \mathcal{S}(\rho_n)$ and $\mu(\Omega) = 1$. In general, $\mathcal{S}(\rho_n) \neq \mathcal{A}$ and μ

may not have a well-behaved extension from \mathcal{A}_0 to all of \mathcal{A} [2, 7]. A subset \mathcal{B} of \mathcal{A} is a *quadratic algebra* if $\emptyset, \Omega \in \mathcal{B}$ and whenever $A, B, C \in \mathcal{B}$ are mutually disjoint with $A \cup B, A \cup C, B \cup C \in \mathcal{B}$, we have $A \cup B \cup C \in \mathcal{B}$. For a quadratic algebra \mathcal{B} , a *q-measure* is a map $\mu_0: \mathcal{B} \rightarrow \mathbb{R}^+$ that satisfies the grade-2 additivity condition (2.1). Of course, an algebra of sets is a quadratic algebra and we conclude that $\mu_n: \mathcal{A}_n \rightarrow \mathbb{R}^+$ is a *q-measure*. It is not hard to show that $\mathcal{S}(\rho_n)$ is a quadratic algebra and $\mu: \mathcal{S}(\rho_n) \rightarrow \mathbb{R}^+$ is a *q-measure* on $\mathcal{S}(\rho_n)$ [3].

3 Classical Sequential Growth Processes

A *partially ordered set (poset)* is a set x together with an irreflexive, transitive relation $<$ on x . In this work we only consider unlabeled posets and isomorphic posets are considered to be identical. Let \mathcal{P}_n be the collection of all posets with cardinality n , $n = 1, 2, \dots$. If $x \in \mathcal{P}_n$, $y \in \mathcal{P}_{n+1}$, then x *produces* y if y is obtained from x by adjoining a single new element to x that is maximal in y . We also say that x is a *producer* of y and y is an *offspring* of x . If x produces y we write $x \rightarrow y$. We denote the set of offspring of x by $x \rightarrow$ and for $A \subseteq \mathcal{P}_n$ we use the notation

$$A \rightarrow = \{y \in \mathcal{P}_{n+1} : x \rightarrow y, x \in A\}$$

The transitive closure of \rightarrow makes the set of all finite posets $\mathcal{P} = \cup \mathcal{P}_n$ into a poset.

A *path* in \mathcal{P} is a string (sequence) x_1, x_2, \dots where $x_i \in \mathcal{P}_i$ and $x_i \rightarrow x_{i+1}$, $i = 1, 2, \dots$. An *n-path* in \mathcal{P} is a finite string $x_1 x_2 \cdots x_n$ where again $x_i \in \mathcal{P}_i$ and $x_i \rightarrow x_{i+1}$. We denote the set of paths by Ω and the set of *n-paths* by Ω_n . The set of paths whose initial *n-path* is $\omega_0 \in \Omega_n$ is denoted by $\omega_0 \Rightarrow$. Thus, if $\omega_0 = x_1 x_2 \cdots x_n$ then

$$\omega_0 \Rightarrow = \{\omega \in \Omega : \omega = x_1, x_2 \cdots x_n y_{n+1} y_{n+2} \cdots\}$$

If x produces y in r isomorphic ways, we say that the *multiplicity* of $x \rightarrow y$ is r and write $m(x \rightarrow y) = r$. For example, in Figure 1, $m(x \rightarrow y) = 3$. (To be precise, these different isomorphic ways require a labeling of the posets and this is the only place that labeling needs to be mentioned.)




Figure 1

If $x \in \mathcal{P}$ and $a, b \in x$ we say that a is an *ancestor* of b and b is a *successor* of a if $a < b$. We say that a is a *parent* of b and b is a *child* of a if $a < b$ and there is no $c \in x$ such that $a < c < b$. Let $c = (c_0, c_1, \dots)$ be a sequence of nonnegative numbers called *coupling constants* [5, 9]. For $r, s \in \mathbb{N}$ with $r \leq s$, we define

$$\lambda_c(s, r) = \sum_{k=r}^s \binom{s-r}{k-r} c_k = \sum_{k=0}^{s-r} \binom{s-r}{k} c_{r+k}$$

For $x \in \mathcal{P}_n$ $y \in \mathcal{P}_{n+1}$ with $x \rightarrow y$ we define the *transition probability*

$$p_c(x \rightarrow y) = m(x \rightarrow y) \frac{\lambda_c(\alpha, \pi)}{\lambda_c(n, 0)}$$

where α is the number of ancestors and π the number of parents of the adjoined maximal element in y that produces y from x . It is shown in [5, 9] that $p_c(x \rightarrow y)$ is a probability distribution in that it satisfies the Markov-sum rule

$$\sum \{p_c(x \rightarrow y) : y \in \mathcal{P}_{n+1}, x \rightarrow y\} = 1$$

In discrete quantum gravity, the elements of \mathcal{P} are thought of as causal sets and $a < b$ is interpreted as b being in the causal future of a . The distribution $y \mapsto p_c(x \rightarrow y)$ is essentially the most general that is consistent with principles of causality and covariance [5, 9]. It is hoped that other theoretical principles or experimental data will determine the coupling constants. One suggestion is to take $c_k = 1/k!$ [6, 7]. The case $c_k = c^k$ for some $c > 0$ has been previously studied and is called a *percolation dynamics* [5, 6, 8].

We call an element $x \in \mathcal{P}$ a *site* and we sometimes call an n -path an *n-universe* and a path a *universe*. The set \mathcal{P} together with the set of transition probabilities $p_c(x \rightarrow y)$ forms a *classical sequential growth process* (CSGP)

which we denote by (\mathcal{P}, p_c) [4, 5, 6, 8, 9]. It is clear that (\mathcal{P}, p_c) is a Markov chain and as usual we define the probability of an n -path $\omega = y_1 y_2 \cdots y_n$ by

$$p_c^n(\omega) = p_c(y_1 \rightarrow y_2) p_c(y_2 \rightarrow y_3) \cdots p_c(y_{n-1} \rightarrow y_n)$$

Denoting the power set of Ω_n by 2^{Ω_n} , $(\Omega_n, 2^{\Omega_n}, p_c^n)$ becomes a probability space where

$$p_c^n(A) = \sum \{p_c^n(\omega) : \omega \in A\}$$

for all $A \in 2^{\Omega_n}$. The probability of a site $x \in \mathcal{P}_n$ is

$$p_c^n(x) = \sum \{p_c^n(\omega) : \omega \in \Omega_n, \omega \text{ ends at } x\}$$

Of course, $x \mapsto p_c^n(x)$ is a probability measure on \mathcal{P}_n and we have

$$\sum_{x \in \mathcal{P}_n} p_c^n(x) = 1$$

Example 1. Figure 2 illustrates the first two steps of a CSGP where the 2 indicates the multiplicity $m(x_3 \rightarrow x_6) = 2$. Table 1 lists the probabilities of the various sites for the general coupling constants c_k and the particular coupling constants $c'_k = 1/k!$ where $d = (c_0 + c_1)(c_0 + 2c_1 + c_2)$.




Figure 2

x_i	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_8
$p_c^{(n)}(x_i)$	1	$\frac{c_1}{c_0+c_1}$	$\frac{c_0}{c_0+c_1}$	$\frac{c_1(c_1+c_2)}{d}$	$\frac{c_1^2}{d}$	$\frac{3c_0c_1}{d}$	$\frac{c_0c_2}{d}$	$\frac{c_0^2}{d}$
$p_c^n(x_i)$	1	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{14}$	$\frac{1}{7}$	$\frac{3}{7}$	$\frac{1}{14}$	$\frac{1}{7}$

Table 1

For $A \subseteq \Omega_n$ we use the notation

$$A \Rightarrow = \cup \{\omega \Rightarrow : \omega \in A\}$$

Thus, $A \Rightarrow$ is the set of paths whose initial n -paths are elements of A . We call $A \Rightarrow$ a *cylinder set* and define

$$\mathcal{A}_n = \{A \Rightarrow : A \subseteq \Omega_n\}$$

In particular, if $\omega \in \Omega_n$ then the *elementary cylinder set* $\text{cyl}(\omega)$ is given by $\text{cyl}(\omega) = \omega \Rightarrow$. It is easy to check that the \mathcal{A}_n form an increasing sequence $\mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$ of algebras on Ω and hence $\mathcal{C}(\Omega) = \cup \mathcal{A}_n$ is an algebra of subsets of Ω . Also for $A \in \mathcal{C}(\Omega)$ of the form $A = A_1 \Rightarrow$, $A_1 \subseteq \Omega_n$, we define $p_c(A) = p_c^n(A_1)$. It is easy to check that p_c is a well-defined probability measure on $\mathcal{C}(\Omega)$. It follows from the Kolmogorov extension theorem that p_c has a unique extension to a probability measure ν_c on the σ -algebra \mathcal{A} generated by $\mathcal{C}(\Omega)$. We conclude that $(\Omega, \mathcal{A}, \nu_c)$ is a probability space, the increasing sequence of subalgebras \mathcal{A}_n generates \mathcal{A} and that the restriction $\nu_c | \mathcal{A}_n = p_c^n$. Hence, the subspaces $H_n = L_2(\Omega, \mathcal{A}_n, p_c^n)$ form a filtration of the Hilbert space $H = L_2(\Omega, \mathcal{A}, \nu_c)$.

4 Quantum Sequential Growth Processes

This section employs the framework of Section 2 to obtain a quantum sequential growth process (QSGP) from the CSGP (\mathcal{P}, p_c) developed in Section 3. We have seen that the *n-path Hilbert space* $H_n = L_2(\Omega, \mathcal{A}_n, p_c^n)$ forms a filtration of the *path Hilbert space* $H = L_2(\Omega, \mathcal{A}, \nu_c)$. In the sequel, we assume that $p_c^n(\omega) \neq 0$ for every $\omega \in \Omega_n$, $n = 1, 2, \dots$. Then the set of vectors

$$e_\omega^n = p_c^n(\omega)^{1/2} \chi_{\text{cyl}(\omega)}, \omega \in \Omega_n$$

form an orthonormal basis for H_n , $n = 1, 2, \dots$. For $A \in \mathcal{A}_n$, notice that $\chi_A \in H$ with $\|\chi_A\| = p_c^n(A)^{1/2}$.

We call a DQP $\rho_n \in \mathcal{Q}(H_n)$ a *quantum sequential growth process* (QSGP). We call ρ_n the *local operators* and $\mu_n(A) = D_n(A, A)$ the *local q -measures* for the process. If $\rho = \lim \rho_n$ exists in the strong operator topology, then ρ is a q -probability operator on H called the *global operator* for the process. If the global operator ρ exists, then $\hat{\mu}(A) = \langle \rho \chi_A, \chi_A \rangle$ is a (continuous) q -measure on \mathcal{A} that extends μ_n , $n = 1, 2, \dots$. Unfortunately, the global operator does not exist, in general, so we must be content to work with the local operators [2, 3, 7]. In this case, we still have the q -measure μ on the quadratic algebra $\mathcal{S}(\rho_n) \subseteq \mathcal{A}$ that extends μ_n $n = 1, 2, \dots$. We frequently identify a set $A \subseteq \Omega_n$ with the corresponding cylinder set $(A \Rightarrow) \in \mathcal{A}_n$. We then have the q -measure, also denoted by μ_n , on 2^{Ω_n} defined by $\mu_n(A) = \mu_n(A \Rightarrow)$. Moreover, we define the q -measure, again denoted by μ_n , on \mathcal{P}_n by

$$\mu_n(A) = \mu_n(\{\omega \in \Omega_n : \omega \text{ end in } A\})$$

for all $A \subseteq \mathcal{P}_n$. In particular, for $x \in \mathcal{P}_n$ we have

$$\mu_n(\{x\}) = \mu_n(\{\omega \in \Omega_n : \omega \text{ ends with } x\})$$

If $A \in \mathcal{A}_n$ has the form $A_1 \Rightarrow$ for $A_1 \subseteq \Omega_n$ then $A \in \mathcal{A}_{n+1}$ and $A = (A_1 \rightarrow) \Rightarrow$ where $A_1 \rightarrow \subseteq \Omega_{n+1}$. Let $\rho_n \in \mathcal{Q}(H_n)$, $\rho_{n+1} \in \mathcal{Q}(H_{n+1})$ and let $D_n(A, B) = \langle \rho_n \chi_B, \chi_A \rangle$, $D_{n+1}(A, B) = \langle \rho_{n+1} \chi_B, \chi_A \rangle$ be the corresponding decoherence functionals. Then ρ_{n+1} is consistent with ρ_n if and only if for all $A, B \subseteq \Omega_n$ we have

$$D_{n+1}[(A \rightarrow) \Rightarrow, (B \rightarrow) \Rightarrow] = D_n(A \Rightarrow, B \Rightarrow) \quad (4.1)$$

Lemma 4.1. *For $\rho_n \in \mathcal{Q}(H_n)$, $\rho_{n+1} \in \mathcal{Q}(H_{n+1})$ we have that ρ_{n+1} is consistent with ρ_n if and only if for all $\omega, \omega' \in \Omega_n$ we have*

$$D_{n+1}[(\omega \rightarrow) \Rightarrow, (\omega' \rightarrow) \Rightarrow] = D_n(\omega \Rightarrow, \omega' \Rightarrow) \quad (4.2)$$

Proof. Necessity is clear. For sufficiency, suppose (4.2) holds. Then for every $A, B \subseteq \Omega_n$ we have

$$\begin{aligned} D_{n+1}[(A \rightarrow) \Rightarrow, (B \rightarrow) \Rightarrow] &= \sum_{\omega \in A} \sum_{\omega' \in B} D_{n+1} D_{n+1}[(\omega \rightarrow) \Rightarrow, (\omega' \rightarrow) \Rightarrow] \\ &= \sum_{\omega \in A} \sum_{\omega' \in B} D_n(\omega \Rightarrow, \omega' \Rightarrow) = D_n(A \Rightarrow, B \Rightarrow) \end{aligned}$$

and the result follows from (4.1). \square

For $\omega = x_1x_2 \cdots x_n \in \Omega_n$ and $y \in \mathcal{P}_{n+1}$ with $x_n \rightarrow y$ we use the notation $\omega y \in \Omega_{n+1}$ where $\omega y = x_1x_2 \cdots x_ny$. We also define $p_c(\omega \rightarrow y) = p_c(x_n \rightarrow y)$ and write $\omega \rightarrow y$ whenever $x_n \rightarrow y$.

Theorem 4.2. *For $\rho_n \in \mathcal{Q}(H_n)$, $\rho_{n+1} \in \mathcal{Q}(H_{n+1})$ we have that ρ_{n+1} is consistent with ρ_n if and only if for every $\omega, \omega' \in \Omega_n$ we have*

$$\langle \rho_n e_{\omega'}^n, e_{\omega}^n \rangle = \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x}} \sum_{\substack{y \in \mathcal{P}_{n+1} \\ \omega \rightarrow y}} p_c(\omega' \rightarrow x)^{1/2} p_c(\omega \rightarrow y)^{1/2} \langle \rho_{n+1} e_{\omega'x}^{n+1}, e_{\omega y}^{n+1} \rangle \quad (4.3)$$

Proof. By Lemma 4.1, ρ_{n+1} is consistent with ρ_n if and only if (4.2) holds. But

$$\begin{aligned} D_n(\omega \Rightarrow, \omega' \Rightarrow) &= \langle \rho_n \chi_{\omega' \Rightarrow}, \chi_{\omega \Rightarrow} \rangle = \langle \rho_n \chi_{\text{cyl}(\omega')}, \chi_{\text{cyl}(\omega)} \rangle \\ &= p_c^n(\omega')^{1/2} p_c^n(\omega)^{1/2} \langle \rho_n e_{\omega'}^n, e_{\omega}^n \rangle \end{aligned}$$

Moreover, we have

$$\begin{aligned} D_{n+1}[(\omega \rightarrow) \Rightarrow, (\omega' \rightarrow) \Rightarrow] &= \langle \rho_{n+1} \chi_{(\omega \rightarrow) \Rightarrow}, \chi_{(\omega' \rightarrow) \Rightarrow} \rangle \\ &= \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x}} \sum_{\substack{y \in \mathcal{P}_{n+1} \\ \omega \rightarrow y}} \langle \rho_{n+1} \chi_{\omega'x \Rightarrow}, \chi_{\omega y \Rightarrow} \rangle \\ &= \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x}} \sum_{\substack{y \in \mathcal{P}_{n+1} \\ \omega \rightarrow y}} \langle \rho_{n+1} \chi_{\text{cyl}(\omega'x)}, \chi_{\text{cyl}(\omega y)} \rangle \\ &= \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x}} \sum_{\substack{y \in \mathcal{P}_{n+1} \\ \omega \rightarrow y}} p_c^n(\omega'x)^{1/2} p_c^n(\omega y)^{1/2} \langle \rho_{n+1} e_{\omega'x}^{n+1}, e_{\omega y}^{n+1} \rangle \\ &= p_c^n(\omega')^{1/2} p_c^n(\omega)^{1/2} \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x}} \sum_{\substack{y \in \mathcal{P}_{n+1} \\ \omega \rightarrow y}} p_c(\omega' \rightarrow x) p_c(\omega \rightarrow y)^{1/2} \langle \rho_n e_{\omega'x}^{n+1}, e_{\omega y}^{n+1} \rangle \end{aligned}$$

The result now follows. \square

Viewing H_n as $L_2(\Omega_n, 2^{\Omega_n}, p_c^n)$ we can write (4.3) in the simple form

$$\langle \rho_n \chi_{\{\omega'\}}, \chi_{\{\omega\}} \rangle = \langle \rho_{n+1} \chi_{\omega' \rightarrow}, \chi_{\omega \rightarrow} \rangle \quad (4.4)$$

Corollary 4.3. *A sequence $\rho_n \in \mathcal{Q}(H_n)$ is a QSGP if and only if (4.3) or (4.4) hold for every $\omega, \omega' \in \Omega_n$, $n = 1, 2, \dots$.*

We now consider pure q -probability operators. In the following results we again view H_n as $L_2(\Omega_n, 2^{\Omega_n}, p_c^n)$.

Corollary 4.4. *If $\rho_n \in \mathcal{Q}_p(H_n)$, $\rho_{n+1} \in \mathcal{Q}_p(H_{n+1})$ with $p_n = |\psi_n\rangle\langle\psi_n|$, $\rho_{n+1} = |\psi_{n+1}\rangle\langle\psi_{n+1}|$, then ρ_{n+1} is consistent with ρ_n if and only if for every $\omega, \omega' \in \Omega_n$ we have*

$$\langle\psi_n, \chi_{\{\omega\}}\rangle\langle\chi_{\{\omega'\}}, \psi_n\rangle = \langle\psi_{n+1}, \chi_{\omega\rightarrow}\rangle\langle\chi_{\omega'\rightarrow}, \psi_{n+1}\rangle \quad (4.5)$$

Corollary 4.5. *A sequence $|\psi_n\rangle\langle\psi_n| \in \mathcal{Q}_p(H_n)$ is a QSGP if and only if (4.5) holds for every $\omega, \omega' \in \Omega_n$.*

We say that $\psi_{n+1} \in \mathcal{V}(H_{n+1})$ is *strongly consistent* with $\psi_n \in \mathcal{V}(H_n)$ if for every $\omega \in \Omega_n$ we have

$$\langle\psi_n, \chi_{\{\omega\}}\rangle = \langle\psi_{n+1}, \chi_{\omega\rightarrow}\rangle \quad (4.6)$$

By (4.5) strong consistency implies the consistency of the corresponding q -probability operators.

Corollary 4.6. *If $\psi_{n+1} \in \mathcal{V}(H_{n+1})$ is strongly consistent with $\psi_n \in \mathcal{V}(H_n)$, $n = 1, 2, \dots$, then $|\psi_n\rangle\langle\psi_n| \in \mathcal{Q}_p(H_n)$ is a QSGP.*

Lemma 4.7. *If $\psi_n \in \mathcal{V}(H_n)$ and $\psi_{n+1} \in H_{n+1}$ satisfies (4.6) for every $\omega \in \Omega_n$, then $\psi_{n+1} \in \mathcal{V}(H_{n+1})$.*

Proof. Since $\psi_n \in \mathcal{V}(H_n)$ we have by (4.6) that

$$\begin{aligned} |\langle\psi_{n+1}, 1\rangle| &= \left| \sum_{\omega \in \Omega_n} \langle\psi_{n+1}, \chi_{\omega\rightarrow}\rangle \right| = \left| \sum_{\omega \in \Omega_n} \langle\psi_n, \chi_{\{\omega\}}\rangle \right| \\ &= |\langle\psi_n, 1\rangle| = 1 \end{aligned}$$

The result now follows. \square

Corollary 4.8. *If $\|\psi_1\| = 1$ and $\psi_n \in H_n$ satisfies (4.6) for all $\omega \in \Omega_n$, $n = 1, 2, \dots$, then $|\psi_n\rangle\langle\psi_n|$ is a QSGP.*

Proof. Since $\|\psi_1\| = 1$, it follows that $\psi_1 \in \mathcal{V}(H_1)$. By Lemma 4.7, $\psi_n \in \mathcal{V}(H_n)$, $n = 1, 2, \dots$. Since (4.6) holds, the result follows from Corollary 4.6. \square

Another way of writing (4.6) is

$$\sum_{\omega \rightarrow x} p_c^{n+1}(\omega x) \psi_{n+1}(\omega x) = p_c^n(\omega) \psi_n(x) \quad (4.7)$$

for every $\omega \in \Omega_n$.

5 Discrete Quantum Gravity Models

This section gives some examples of QSGP that can serve as models for discrete quantum gravity. The simplest way to construct a QSGP is to form the constant pure DQP $\rho_n = |1\rangle\langle 1|$, $n = 1, 2, \dots$. To show that ρ_n is indeed consistent, we have for $\omega \in \Omega_n$ that

$$\sum_{\omega \rightarrow x} p_c^{n+1}(\omega x) = \sum_{\omega \rightarrow x} p_c^n(\omega) p_c(\omega \rightarrow x) = p_c^n(\omega) \sum_{\omega \rightarrow x} p_c(\omega \rightarrow x) = p_c^n(\omega)$$

so consistency follows from (4.7). The corresponding q -measures are given by

$$\mu_n(A) = |\langle 1, \chi_A \rangle|^2 = p_c^n(A)^2$$

for every $A \in \mathcal{A}_n$. Hence, μ_n is the square of the classical measure. Of course, $|1\rangle\langle 1|$ is the global q -probability operator for this QSGP and in this case $\mathcal{S}(\rho_n) = \mathcal{A}$. Moreover, we have the global q -measure $\mu(A) = \nu_c(A)^2$ for $A \in \mathcal{A}$.

Another simple way to construct a QSGP is to employ Corollary 4.8. In this way we can let $\psi_1 = 1$, ψ_2 any vector in $L_2(\Omega_2, 2^{\Omega_2}, p_c^2)$ satisfying

$$\langle \psi_2, \chi_{\{x_1 x_2\}} \rangle + \langle \psi_2, \chi_{\{x_1 x_3\}} \rangle = \langle \psi_1, \chi_{\{x_1\}} \rangle = 1$$

and so on, where x_1, x_2, x_3 are given in Figure 2. As a concrete example, let $\psi_1 = 1$,

$$\psi_2 = \frac{1}{2} [p_c^2(x_1 x_2)^{-1} \chi_{\{x_1 x_2\}} + p_c^2(x_1 x_3) \chi_{\{x_1 x_3\}}]$$

and in general

$$\psi_n = \frac{1}{|\Omega_n|} \sum_{\omega \in \Omega_n} p_c^n(\omega)^{-1} \chi_{\{\omega\}}$$

The q -measure μ_1 is $\mu_1(\{x_1\}) = 1$ and μ_2 is given by

$$\begin{aligned} \mu_2(\{x_1 x_2\}) &= |\langle \psi_2, \chi_{\{x_1 x_2\}} \rangle|^2 = \frac{1}{4} \\ \mu_2(\{x_1 x_3\}) &= |\langle \psi_2, \chi_{\{x_1 x_3\}} \rangle|^2 = \frac{1}{4} \\ \mu_2(\Omega_2) &= |\langle \psi_2, 1 \rangle|^2 = 1 \end{aligned}$$

In general, we have $\mu_n(A) = |A|^2 / |\Omega_n|^2$ for all $A \in \Omega_n$. Thus μ_n is the square of the uniform distribution. The global operator does not exist because there is no q -measure on \mathcal{A} that extends μ_n for all $n \in \mathbb{N}$. For $A \in \mathcal{A}$ we have

$$\langle \psi_n, \chi_A \rangle = \int \psi_n \chi_A d\nu_c = \frac{|A \cap \{\text{cyl}(\omega) : \omega \in \Omega_n\}|}{|\Omega_n|}$$

Letting $\rho_n = |\psi_n\rangle\langle\psi_n|$ we conclude that $A \in \mathcal{S}(\rho_n)$ if and only if

$$\lim_{n \rightarrow \infty} \frac{|A \cap \{\text{cyl}(\omega) : \omega \in \Omega_n\}|}{|\Omega_n|}$$

exists. For example, if $|A| < \infty$ then for n sufficiently large we have

$$|A \cap \{\text{cyl}(\omega) : \omega \in \Omega_n\}| = |A|$$

so $A \in \mathcal{S}(\rho_n)$ and $\mu(A) = 0$. In a similar way if $|A| < \infty$ then for the complement A' , if n is sufficiently large we have

$$|A' \cap \{\text{cyl}(\omega) : \omega \in \Omega_n\}| = |\Omega_n| - |A|$$

so $A' \in \mathcal{S}(\rho_n)$ with $\mu(A') = 1$.

We now present another method for constructing a QSGP. Unlike the previous method this DQP is not pure. Let $\alpha_\omega \in \mathbb{C}$, $\omega \in \Omega_n$ satisfy

$$\left| \sum_{\omega \in \Omega_n} \alpha_\omega p_c^n(\omega)^{1/2} \right| = 1 \quad (5.1)$$

and let ρ_n be the operator on H_n satisfying

$$\langle \rho_n e_\omega^n, e_{\omega'}^n \rangle = \alpha_{\omega'} \overline{\alpha_\omega} \quad (5.2)$$

Then ρ_n is a positive operator and by (5.1), (5.2) we have

$$\begin{aligned} \langle \rho_n 1, 1 \rangle &= \left\langle \rho_n \sum_{\omega} p_c^n(\omega)^{1/2} e_\omega^n, \sum_{\omega'} p_c^n(\omega')^{1/2} e_{\omega'}^n \right\rangle \\ &= \sum_{\omega, \omega'} p_c^n(\omega)^{1/2} p_c^n(\omega')^{1/2} \langle \rho_n e_\omega^n, e_{\omega'}^n \rangle \\ &= \left| \sum_{\omega} p_c^n(\omega)^{1/2} \alpha_\omega \right|^2 = 1 \end{aligned}$$

Hence, $\rho_n \in \mathcal{Q}(H_n)$. Now

$$\Omega_{n+1} = \{\omega x : \omega \in \Omega_n, x \in \mathcal{P}_{n+1}, \omega \rightarrow x\}$$

and for each $\omega x \in \Omega_{n+1}$, let $\beta_{\omega x} \in \mathbb{C}$ satisfy

$$\left| \sum_{\omega x \in \Omega_{n+1}} \beta_{\omega x} p_c^{n+1}(\omega x)^{1/2} \right| = 1$$

Let ρ_{n+1} be the operator on H_{n+1} satisfying

$$\langle \rho_{n+1} e_{\omega x}^{n+1}, e_{\omega' x'}^{n+1} \rangle = \beta_{\omega' x'} \overline{\beta_{\omega x}} \quad (5.3)$$

As before, we have that $\rho_{n+1} \in \mathcal{Q}(H_{n+1})$. The next result follows from Theorem 4.2.

Theorem 5.1. *The operator ρ_{n+1} is consistent with ρ_n if and only if for every $\omega, \omega' \in \Omega_n$ we have*

$$\alpha_{\omega'} \overline{\alpha_{\omega}} = \sum_{\substack{x' \in \mathcal{P}_{n+1} \\ \omega' \rightarrow x'}} \beta_{\omega' x'} p_c(\omega' \rightarrow x')^{1/2} \sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega \rightarrow x}} \overline{\beta_{\omega x}} p_c(\omega \rightarrow x)^{1/2} \quad (5.4)$$

A sufficient condition for (5.4) to hold is

$$\sum_{\substack{x \in \mathcal{P}_{n+1} \\ \omega \rightarrow x}} \beta_{\omega x} p_c(\omega \rightarrow x)^{1/2} = \alpha_{\omega} \quad (5.5)$$

The proof of the next result is similar to the proof of Lemma 4.7.

Lemma 5.2. *Let $\rho_n \in \mathcal{Q}(H_n)$ be defined by (5.2) and let ρ_{n+1} be the operator on H_{n+1} defined by (5.3). If (5.5) holds, then $\rho_{n+1} \in \mathcal{Q}(H_{n+1})$ and ρ_{n+1} is consistent with ρ_n .*

The next result gives the general construction.

Corollary 5.3. *Let $\rho_1 = I \in \mathcal{Q}(H_1)$ and define $\rho_n \in \mathcal{Q}(H_n)$ inductively by (5.3). Then ρ_n is a QSGP.*

References

- [1] L. Bombelli, J. Lee, D. Meyer and R. Sorkin, Spacetime as a casual set, *Phys. Rev. Lett.* **59** (1987), 521–524.

- [2] F Dowker, S. Johnston and S. Surya, On extending the quantum measure, arXiv: quant-ph 1002:2725 (2010).
- [3] S. Gudder, Discrete quantum gravity, arXiv: gr-qc 1108.2296 (2011).
- [4] J. Henson, Quantum histories and quantum gravity, arXiv: gr-qc 0901.4009 (2009).
- [5] D. Rideout and R. Sorkin, A classical sequential growth dynamics for causal sets, *Phys. Rev. D* **61** (2000), 024002.
- [6] R. Sorkin, Causal sets: discrete gravity, arXiv: gr-qc 0309009 (2003).
- [7] R. Sorkin, Toward a “fundamental theorem of quantal measure theory,” arXiv: hep-th 1104.0997 (2011) and *Math. Struct. Comm. Sci.* (to appear).
- [8] S. Surya, Directions in causal set quantum gravity, arXiv: gr-qc 1103.6272 (2011).
- [9] M. Varadarajan and D. Rideout, A general solution for classical sequential growth dynamics of causal sets, *Phys. Rev. D* **73** (2006), 104021.